

Improving water-use efficiency under fluctuating light

Charles P. Pignon¹ * (pignon2@illinois.edu), Samuel Bonfim-Fernandes¹, Ravi Valluru², Nonoy Bandillo², Michael Gore², Edward Buckler², Ivan Baxter³, Stephen P. Long¹, and **Andrew D. B. Leakey**¹.

¹University of Illinois at Urbana-Champaign, Urbana, IL;

²Cornell University, Ithaca, NY;

³Danforth Center, St. Louis, MO

www.foxmillet.org

Project Goals:

This project aims to leverage *Setaria viridis* as a model system to develop novel technologies and methodologies to redesign the bioenergy feedstock *Sorghum bicolor* to enhance water use and photosynthetic efficiencies.

Abstract

In leaves, efficient water-use requires coordination of photosynthetic CO₂ assimilation with the activity of stomata, specialized cellular complexes which regulate CO₂ and water fluxes between the leaf and atmosphere. Stomata are slower than photosynthesis to respond to changes in light, and poor coordination between stomatal and photosynthetic processes can diminish productivity and water-use efficiency. Therefore, acceleration of stomatal responses has been hypothesized to improve water-use efficiency in fluctuating light conditions, which predominate in field crop canopies.

We used high-throughput thermal imaging to assess the speed of stomatal closure following a drop in light in a diversity panel of 667 accessions of *Sorghum bicolor* L. Different accessions showed various speeds of initial stomatal closure after the light was dimmed, and varying responses of stomatal re-opening at low light. The initial speed of stomatal closure, and the level of stomatal conductance at low light, were moderately heritable ($h_g > 0.45$). Mapping with SNPs and gene expression data associated stomata kinetic traits to several candidate genes.

Funding statement.

This project is funded by grant DE-SC0018277 from The DOE Department of Biological and Environmental Research